

# West Central Veterinary Services

---

316 N Jackson St  
Greencastle, IN 46135  
Phone: (765) 301-4369

1472 S US Hwy 41  
Rockville, IN 47872  
Phone: (765) 569-3210

212 W US Hwy 136  
Veedersburg, IN 47987  
Phone: (765) 294-2203

---

## Internal Parasites in Goats

### Common parasites

#### Trichostrongyles

***Haemonchus contortus*** (barber pole worm) → Lives in the abomasum (part of stomach) and causes anemia (pale mucous membranes), protein loss (bottle jaw), weakness, weight loss, and death. One adult female can produce 5,000 to 10,000 eggs in a day! This is the most important parasite because it has developed resistance to multiple dewormers.

***Trichostrongylus*** (bankrupt worm), ***Teladorsagia*** (brown stomach worm), ***Nematodirus***, ***Cooperia*** → Causes diarrhea, protein loss (bottle jaw), and weight loss.

#### Hookworms

***Bunostomum*** → Causes anemia, protein loss, and weight loss; not seen nearly as often as the trichostrongyles.

#### Tapeworms

***Moniezia***, ***Thysanosoma*** "fringed tapeworm" → Only causes problems such as poor weight gain if animal has a heavy worm burden. May see the proglottids (segments) of the worm on rear and tail.

#### Threadworms

***Strongyloides papillosus*** → Rarely causes problems, except in neonates.

#### Whipworms

***Trichuris ovis*** → Rarely causes problems, except if young have heavy load of this parasite. In those cases it causes diarrhea and weight loss.

#### Nodular worm

***Oesophagostomum sp.*** → Causes severe inflammation in the large bowel walls resulting in diarrhea, protein loss, weight loss, lack of appetite, and weakness.

#### Liver Flukes


***Fasciola hepatica***, ***Fascioloides magna*** → Causes severe inflammation in the liver resulting in lethargy, lack of appetite, weight loss, and possibly sudden death. This is not seen very often in this region.

#### Coccidia

***Eimeria sp.*** → Causes diarrhea, dehydration, weight loss, death. Clinical signs are worse in young goats. Overcrowding allows for excessive manure build-up which is ideal for the propagation of coccidia.

\*\*Treatment and control of coccidia will be addressed in a separate section from the other parasites

## Life Cycle of *Haemonchus contortus*:



While the life cycle of other parasites is important, *Haemonchus* is our primary concern, so it is useful to understand its life cycle when developing a control program. The adult worm lives in the abomasum (one of the stomachs) where they attach and suck blood. Eggs are laid and are then passed in the feces. Eggs hatch into the 1<sup>st</sup> stage larvae and then mature to 2<sup>nd</sup> and 3<sup>rd</sup> stage larvae in the pasture. 3<sup>rd</sup> stage larvae migrate up blades of grass where the goats ingest them as they graze. The larvae mature to adults in the goat and the cycle begins again.

## Control of Parasites:

Depending on the number of animals, acreage/lot size, and history of parasite control, deworming recommendations will vary from farm to farm. Resistance to dewormers has become a major problem in goats. *Haemonchus*, in particular, has become resistant to many dewormers in various regions of the U.S. It is important to keep in mind that parasite control involves more than just deworming your goats.

- The goal in controlling parasites should not be to get rid of every single parasite but to **decrease the population of parasites** so that they do not cause serious herd problems.
- **Minimize fecal contamination of feeds.** Don't place feed on the ground and keep feed troughs cleaned out.
- **Deworming too often is one of the major causes of parasite resistance to dewormers.**
  - When questioning the efficacy of a dewormer, it is helpful to get a fecal sample prior to deworming and then collect a second sample 2-3 weeks after deworming. We can perform a fecal egg count. If the dewormer is effective there should be a 90-95% reduction in the number of eggs.
- **Isolate** new animals that are brought onto the farm. These animals need to be dewormed and then have the feces checked for parasitic eggs 2 weeks after deworming. Once they are clear of parasites, they can be mixed with the rest of the herd. Isolation is a good practice to follow to prevent other diseases as well.
- Be sure to **use the appropriate dose of dewormer**. Underdosing is not only ineffective, it can also lead to resistant parasites.
- **Pasture management- "The solution to pollution is dilution."**
  - This phrase holds true for parasite control in any species. Parasites are going to be a bigger problem when you have lots of animals in a small area. The ideal stocking rate for goats (or sheep) is **6-8 animals per acre of land**.

- Most of the worm burden is found in the pasture. Here are a couple of ways to decrease the amount of larvae in the pasture:
  - There will be some larvae that will survive the winter in the pasture. About 3 weeks after the animals have been on the spring pasture, deworm them all. This will kill the worms that the goats have gotten from ingesting the larvae that survived the winter. This may need to be repeated 1-2 more times 3-4 weeks apart.
  - If you have access to multiple pastures, you may be able to use a pasture for weeks to months before the larvae build up on the pasture. Then you can deworm the herd and rotate them to a new pasture. This is referred to as a treat and move strategy.
  - Rotating animals on a pasture can also decrease the larvae contamination. Rotate the goats with another animal, like cattle or horses. Do not rotate with sheep because they have the same parasites.
- **FAMACHA**→ This a salvage deworming program. It only works to control *Haemonchus* infestations. It is a process where the color of the inside of the lower eyelid is compared to that on a chart. Those with a pale mucous membrane color are the animals that are treated. The program was developed to minimize the number of animals being dewormed so that resistance was less likely to be a problem. Keep in mind this entails some strategic planning and time to examine each animal. For more information, please visit this website: <http://www.scsr.org>
- **Deworming on a schedule**
  - Scheduled deworming is not the most optimal for preventing the development of resistant parasites; however for many clients, this is the most practical.
  - Keep in mind that these are rough guidelines, we can make adjustments according to your herd's individual needs.
    - Deworm every 3 months.
    - Most of the time use cydectin.
    - Every 3<sup>rd</sup> deworming use either panacur, safeguard, or valbazen rather than cydectin.
    - 2-3 times a year or whenever a parasite problem arises, bring in a few fecal samples **prior** to deworming. Then bring in a few fecal samples **2 weeks after deworming**. This helps us to tailor a deworming program best suited for your herd.

\*\*\*Most research suggests that you should use one dewormer until it is no longer effective. The one problem with that is there are some parasites that may not be killed by a particular dewormer. Our practices have found that a large majority of *Haemonchus* parasites are resistant to ivermectin; however do not discount the benefit of ivermectin for controlling other parasites.

## Common Parasiticides Used in Goats

Drug	Goat Dosage	Meat With-drawal	Milk With-drawal	Haemonchus	Trichostrongylus	Teladorsagia	Nematodirus	Cooperia	Bunostomum	Moneizia	Thysanosoma	Strongyloides	Trichuris	Oesophagostomum	Liver Flukes
<b>Avermectins</b>															
<b>Moxidectin</b> Cydectin drench	3 ml/22 lbs BW (BW-body weight)	7 days	***	X	X	X	X	X	X						X
<b>Ivermectin</b> - Ivomec drench or injectable	Drench- 5 ml/30 lbs BW Inj.- 1 ml/70 lbs BW, give subcutaneously	Drench- 11 days; Inj- 35 days	***	X	X	X	X	X	X			X	X	X	
<b>Doramectin</b> - Dectomax injectable	1 ml/70 lbs BW	??(40 d in cattle)	***	X	X	X	X	X	X			X	X	X	
<b>Eprinomectin</b> - Eprinex- used orally	1 ml/20 lb BW	None	None	X	X	X	X	X	X			X	X	X	
<b>Benzimidazoles</b>															
<b>Albendazole</b> - Valbazen	1 ml/20 lb BW	7 days	***	X	X	X	X	X	X	X	X			X	X
<b>Fenbendazole</b> - Panacur, Safeguard	2 ml/25 lb BW	8 days-cattle	***	X	X	X	X	X	X			X	X	X	
<b>Cholinergics</b>															
<b>Levamisole</b> - Levasol drench	1 ml/25 lb BW	72 hours	***	X	X	X	X	X	X					X	

\*\*\*Milk withdrawal has not been established. We recommend withholding the milk for 96 hours (8 milkings) before using it for human consumption.

Valbazen should not be given to does within the first 30 days of pregnancy.

Levamisole has been ineffective for many farms. If you have not used this drug for many years, it may be useful to try it. It is currently unavailable, but there should be a generic form coming out soon.

Classes of dewormers are listed (avermectins, benzimidazoles, and cholinergics). The main importance is that once parasites become resistant to a dewormer, all the dewormers within that class will be unlikely to work. The exception is Cydectin. Many times Haemonchus may be resistant to Ivomec but are still susceptible to Cydectin.

Pay close attention to the dosages listed above. They are often 2-3 times higher than those listed for sheep.

## Coccidia

Coccidia is a unique protozoal parasite that can be **devastating** in a herd.

- It is a common cause of **pasty to watery diarrhea in kids**. Other signs include dehydration, lack of appetite, weakness, rough hair coat, and death. Disease becomes worse when the goats are stressed (by shipping, weaning, or feed changes).
- Coccidia are transmitted by ingesting feces that contain coccidia.
- The coccidia destroy the cells lining the intestinal tract as they reproduce inside the animal.
- As with the other parasites, "the solution to pollution is dilution". Too many animals in a small area will contribute to the severity of coccidiosis.
- Coccidia do not survive well in dry conditions with direct sunlight.
- Diagnosis of coccidia is by submitting a fecal sample to be examined under a microscope.
- Treatment/Prevention:

<b>Drug</b>	<b>Trade Name</b>	<b>Dose</b>	<b>Comment</b>
Lasalocid	Bovatec	20 30 grams per ton of feed	Used for prevention of coccidia
Decoquinate	Deccox	0.5 lbs per ton of feed	Used for prevention of coccidia
Monensin	Rumensin	15-20 grams per ton of feed	Used for prevention of coccidia
Amprolium	Corid	Drench: 9 mls/50 lbs body weight for 5 days straight	24 hour meat withdrawal; 96 hour milk withdrawal
		In the water: 4 oz of corid per 5 gallons of water. Give as only water source for 5 days straight	